



FAST FACTS HEALTH

One study showed that when a teen announces a transgender identity to their peer group, the number of friends who also became transgender-identified was

3.5

per group.²



In just seven years, there has been a nearly

2,000% increase

in children seeking treatment for sexual identity confusion in the United Kingdom.³



Up to **98%**

of children who struggle with their sex as a boy or a girl, come to accept their sex by adulthood.⁴



Identifying as transgender or nonbinary may be linked to autism spectrum disorders. Children with autism spectrum disorders are **7 times** more likely to want to be the opposite sex than the general population.⁵



After sex reassignment surgery, transgender-identified people are nearly

20 times

more likely to die from suicide than the general population.⁶



Studies show that

100% of children who use puberty blockers will go on to use cross-sex hormones, leaving them permanently sterile.⁷



Girls as young as **13** are undergoing double mastectomies and boys as young as **17** are undergoing full genital sex reassignment surgeries.^{8,9}



The **long-term effects**

of puberty blockers and cross-sex hormones have not been studied.¹⁰



Science demonstrates that there are **two** sex chromosomes—two X chromosomes in females and an X and a Y in males—in nearly every single cell in our bodies.¹¹



Some transgender-identified patients are being prescribed cross-sex hormones on their very **first** visit to a clinic.¹²

